



Labsmart Software

Sample Letterhead

+91 12345 67890

yourlabname@gmail.com

<https://www.yourlabname.in/>**Mr. Dummy**

Age / Sex : 28 YRS / M
Referred by : Dr. Sachin Patil (MBBS)
Reg. no. : 1116



1116

Registered on : 25/11/2024 01:19 PM
Collected on : 25/11/2024
Received on : 25/11/2024
Reported on : 25/11/2024 01:19 PM

Scan to download

**CLINICAL PATHOLOGY**

TEST	VALUE	UNIT	REFERENCE
URINE CORTISOL	11	µg/24 hrs	

Physiological Basis:

Urinary free cortisol measurement is useful in the initial evaluation of suspected Cushing syndrome.

Interpretation**Increased in:**

Cushing syndrome, acute illness, stress.

Not increased in: Obesity.

Comments

Urinary free cortisol is the initial diagnostic test of choice for Cushing syndrome. Not useful for the diagnosis of adrenal insufficiency. A shorter (12-hour) overnight collection and measurement of the ratio of urine-free cortisol to urine creatinine appears to perform nearly as well as a 24-hour collection for urine-free cortisol.

~~~ End of report ~~~

Mr. Sachin Sharma  
DMLT, Lab Incharge

Dr. A. K. Asthana  
MBBS, MD Pathologist

**NOT VALID FOR MEDICO LEGAL PURPOSE**

Work timings: Monday to Sunday, 8 am to 8 pm

Please correlate clinically. Although the test results are checked thoroughly, in case of any unexpected test results which could be due to machine error or typing error or any other reason please contact the lab immediately for a free evaluation.