



Labsmart Software Sample Letterhead

+91 12345 67890

yourlabname@gmail.com

<https://www.yourlabname.in/>**Mr. Dummy**Age / Sex : 28 YRS / M
Referred by : Dr. Sachin Patil (MBBS)
Reg. no. : 1101Registered on : 19/11/2024 06:16 PM
Collected on : 19/11/2024
Received on : 19/11/2024
Reported on : 19/11/2024 06:16 PM

Scan to download



SEROLOGY & IMMUNOLOGY ANTI PHOSPHOLIPID ANTIBODIES

TEST	VALUE	UNIT	REFERENCE
ANTI PHOSPHOLIPID IGG	1	GPL U/mL	
ANTI PHOSPHOLIPID IGM	1	MPL U/mL	

2006 International Consensus statement on Classification of definite APS

CLINICAL CRITERIA	LABORATORY CRITERIA
Arterial/Venous thrombosis	Cardiolipin antibodies (aCL)
Fetal Loss	Beta 2 Glycoprotein 1 antibodies
Premature birth	Lupus Anticoagulant (LA)

Note:

APS is diagnosed if at least one laboratory criterion and one clinical criterion are met. For the laboratory criteria, they need to be observed on two or more separate occasions, with each test done at least 12 weeks apart.

Comments

A phospholipid antibody test measures the levels of autoantibodies against cardiolipin, phosphatidylserine, phosphatidylinositol, and phosphatidic acid. It helps diagnose an increased risk of blood clots in patients with systemic lupus erythematosus (20-35% of cases) and other similar disorders. This test is also used to diagnose Antiphospholipid syndrome (APS) in patients with recent miscarriage (11-22% in all trimesters), Pulmonary hypertension, Non-vegetative endocarditis, Livedo Reticularis, Stroke at young age and Deep vein thrombosis.

~~~ End of report ~~~

Mr. Sachin Sharma  
DMLT, Lab InchargeDr. A. K. Asthana  
MBBS, MD Pathologist

NOT VALID FOR MEDICO LEGAL PURPOSE

Work timings: Monday to Sunday, 8 am to 8 pm

Please correlate clinically. Although the test results are checked thoroughly, in case of any unexpected test results which could be due to machine error or typing error or any other reason please contact the lab immediately for a free evaluation.