

+91 12345 67890

https://www.yourlabname.in/

Mr. Dummy

Age / Sex : 28 YRS / M

Referred by : Dr. Sachin Patil (MBBS)

Reg. no. : 1100

1100

Registered on : 19/11/2024 06:07 PM

Collected on : 19/11/2024 Received on : 19/11/2024

Reported on : 19/11/2024 06:07 PM



SEROLOGY & IMMUNOLOGY BETA 2 GLYCOPROTEIN 1, ANTIBODIES

TEST	VALUE	UNIT	REFERENCE
BETA 2 GLYCOPROTEIN 1, IGG	3	SGU	
BETA 2 GLYCOPROTEIN 1, IGM	3	SMU	

2006 International Consensus statement on Classification of definite APS

CLINICAL CRITERIA	LABORATORY CRITERIA	
Arterial/Venous thrombosis	Cardiolipin antibodies (aCL)	
Fetal Loss	Beta 2 Glycoprotein 1 antibodies	
Premature birth	Lupus Anticoagulant (LA)	

Note:

APS is diagnosed if at least one laboratory criterion and one clinical criterion are met. For the laboratory criteria, they need to be observed on two or more separate occasions, with each test done at least 12 weeks apart.

Comments

This test checks for IgG antibodies to Beta-2 glycoprotein. Finding these antibodies can indicate a risk of blood clots in arteries and veins, as well as a higher chance of repeated miscarriages. The new International classification criteria for APS include Beta-2 glycoprotein for 2 key reasons namely:

- 1. Beta-2 Glycoprotein antibodies maybe the only antibody present in 10% patients of Antiphospholipid syndrome (APS)
- 2. Beta-2 Glycoprotein antibodies are highly specific for APS unlike Anticardiolipin antibodies which may be positive in certain infectious diseases

Indications

- · Clinical history suggestive of Arterial or venous thrombosis
- · History of Recurrent fetal loss /Intrauterine growth restriction/ Premature birth

~~~ End of report ~~~

Mr. Sachin Sharma DMLT, Lab Incharge

Page 1 of 1

**Dr. A. K. Asthana**MBBS, MD Pathologist

# NOT VALID FOR MEDICO LEGAL PURPOSE

Work timings: Monday to Sunday, 8 am to 8 pm