

- +91 12345 67890
- 🖂 yourlabname@gmail.com
- https://www.yourlabname.in/

Mr. Dummy

Age / Sex	: 28 YRS / M
Referred by	: Dr. Sachin Patil (MBBS)
Reg. no.	: 1081





SEROLOGY & IMMUNOLOGY

TEST	VALUE	UNIT	REFERENCE	
VDRL	NON-REACTIV	E		
Result	Remarks			
Reactive	Indicates the presence of IaM & IaG antibodies a	gainst non-treponemal a	antigens	

Reactive	active Indicates the presence of IgM & IgG antibodies against non-treponemal antigens	
Non-Reactive	Indicates absence of IgM & IgG antibodies against non-treponemal antigens	

Note:

1. Titers of \geq 1: 8 and rising titres are significant.

2. Positive result indicates ongoing or recent infection and the diagnosis should be confirmed by specific Treponemal tests such as TPHA & FTA- AbS.

3. The reactivity will vary with the Primary (60-86%), Secondary (99%) and Tertiary (98%) stage of Syphilis.

4. False positive results may be observed in patients of Malaria, Hepatitis, Mumps, Leprosy, Infectious Mononucleosis, Rheumatoid Arthritis and Collagen disease.

5. False negative reaction may be due to processing of sample collected early in the course of disease, immunosuppression and due to prozone effect.

Uses :

- To screen for presence and monitor the progression of Syphilis infection.

- To assess the response to therapy (decreasing titres) in patients being treated for Syphilis.

~~~ End of report ~~~

Saching

Mr. Sachin Sharma DMLT, Lab Incharge

Dr. A. K. Asthana MBBS, MD Pathologist

Page 1 of 1

#### NOT VALID FOR MEDICO LEGAL PURPOSE

Work timings: Monday to Sunday, 8 am to 8 pm

Please correlate clinically. Although the test results are checked thoroughly, in case of any unexpected test results which could be due to machine error or typing error or any other reason please contact the lab immediately for a free evaluation.